

On Locally ϕ -Symmetric β -Kenmotsu Manifolds

ABSOS ALI SHAIKH, SHYAMAL KUMAR HUI

*Department of Mathematics, University of Burdwan, Burdwan – 713104,
West Bengal, India, aask2003@yahoo.co.in*

Presented by Oscar García Prada

Received April 24, 2009

Abstract: The present paper deals with a study of locally ϕ -symmetric β -Kenmotsu manifold with the existence of β -Kenmotsu manifold by an interesting example.

Key words: Conformally flat, weakly locally ϕ -symmetric, strongly locally ϕ -symmetric, β -Kenmotsu, Einstein manifold, scalar curvature.

AMS *Subject Class.* (2000): 53C15, 53C25.

REFERENCES

- [1] D. E. BLAIR, Contact manifolds in Riemannian geometry, *Lecture Notes in Mathematics* 509, Springer-Verlag, Berlin-New York, 1976.
- [2] K. KENMOTSU, A class of almost contact Riemannian manifolds, *Tôhoku Math. J. (2)* **24** (1972), 93–103.
- [3] J. A. OUBIÑA, New classes of almost contact metric structures, *Publ. Math. Debrecen* **32** (3-4) (1985), 187–193.
- [4] A. A. SHAIKH, K. K. BAISHYA, On ϕ -symmetric LP-Sasakian manifolds, *Yokohama Math. J.* **52** (2) (2006), 97–112.
- [5] T. TAKAHASHI, Sasakian ϕ -symmetric spaces, *Tôhoku Math. J. (2)* **29** (1) (1977), 91–113.