

Moufang H^* -algebras

JOSÉ ANTONIO CUENCA MIRA*

*Departamento de Álgebra, Geometría y Topología, Facultad de Ciencias,
Universidad de Málaga, 29080 Málaga, Spain*

(Research paper presented by A. Rodríguez Palacios)

AMS Subject Class. (2000): 17A60, 17D05, 46K70

Received July 30, 2001

We recall that a nonassociative algebra V over \mathbb{K} ($= \mathbb{R}$ or \mathbb{C}) is said to be an H^* -algebra when V is also a Hilbert space over \mathbb{K} (the inner product of which is denoted by $(|)$) and is endowed with an algebra involution \star which is linear in the real case and conjugate-linear in the complex case, and satisfies

$$(xy|z) = (x|zy^\star) = (y|x^\star z)$$

for all $x, y, z \in V$. H^* -algebras were introduced by W. Ambrose [2] in a complex associative setting in a slightly different way. The real associative case was studied by I. Kaplansky [39] (see also [5, 10, 18]). Today the structure theory of several particular classes of nonassociative H^* -algebras is well-known [47, 48, 3, 4, 35, 49, 43, 36, 37, 25, 32, 26, 29, 7, 11, 12, 34, 8, 22, 30, 42, 33]. There is also a complete determination of a wide class of ternary H^* -structures [40, 41, 18, 12, 20, 17, 16, 23, 24, 58], as well as a germinal theory for both binary and ternary arbitrary H^* -structures (see [32, 34, 31, 38, 6, 9, 14, 15, 56, 50, 13, 54, 53, 45] and [19, 21, 57, 51, 52, 58], respectively). The reader is referred to Section E of [44] for a complete survey on H^* -theory.

Let V be an H^* -algebra. Then the left annihilator

$$\text{Lann } V = \{x \in V : xV = 0\}$$

coincides with the right annihilator

$$\text{Rann } V = \{x \in V : Vx = 0\}$$

* This work was partially supported by DGYCIT (PB97-1497), AECI and PAI (FQM-0125).

[32, Proposition 2]. We denote it by $\text{Ann } V$. There is a decomposition of V as an orthogonal direct sum

$$V = \overline{\text{Lin } V^2} \oplus \text{Ann } V, \quad (1)$$

where $\overline{\text{Lin } V^2}$ is the closed linear span of the set $\{xy : x, y \in V\}$ (see [32, Proposition 2] and [34, Theorem 1]). Moreover $\overline{\text{Lin } V^2}$, under a suitable involution and the restriction of the inner product and the algebraic operations of V , is an H^* -algebra with zero annihilator in itself [25, 32, 34]. On the other hand, if V has zero annihilator, then V splits as an orthogonal direct sum

$$V = \bigoplus_{\alpha \in \mathcal{A}} I_{\alpha}, \quad (2)$$

where $\{I_{\alpha}\}$ is the family of all the minimal closed ideals of V , being furthermore every I_{α} a topologically simple H^* -algebra with the structure inherited from V [32, 34, 38]. We recall that an H^* -algebra W is called *topologically simple* if it has nonzero product and its unique closed two-sided ideals are 0 and W .

The decompositions given by (1) and (2) reduce the study of every class of nonassociative H^* -algebras defined by identities to the determination of those H^* -algebras in the class that are topologically simple. This note is devoted to the description of H^* -algebras satisfying some of the Moufang identities. We recall that an *alternative algebra* is a nonassociative algebra satisfying the alternative identities given by

$$x^2y = x(xy), \quad (yx)x = yx^2.$$

It is well known that every alternative algebra satisfies the following identities

$$y((xz)x) = ((yx)z)x \quad (3)$$

$$(xy)(zx) = (x(yz))x \quad (4)$$

$$(x(yx))z = x(y(xz)) \quad (5)$$

(see [55, 46]). These identities are called *right*, *middle*, and *left Moufang* identity, respectively.

In this note we prove that H^* -algebras satisfying some of the Moufang identities are in fact alternative. A similar result on alternativeness of middle Moufang algebras has been recently obtained in [27] and [28] for division and composition middle Moufang algebras. Applying among other results a deep theorem of M. Slater [55, Theorem 9, p. 194], alternative H^* -algebras can be completely described [44, p. 148] (see also [43]). It follows that the result in this note concludes the structure theory of Moufang H^* -algebras.

1. THE RESULTS

Let A be a nonassociative algebra over a field F and $g : A \times A \rightarrow F$ a symmetric bilinear form. We recall that g is said to be associative if

$$g(xy, z) = g(x, yz)$$

for all $x, y, z \in A$.

LEMMA 1.1. *Let A be a nonassociative algebra over a field F such that there exists a nondegenerate associative symmetric bilinear form on A . Then the following assertions are equivalent:*

- (i) *A satisfies the right Moufang identity.*
- (ii) *A satisfies the middle Moufang identity.*
- (iii) *For all $x, z, u \in A$ the equality $((xz)x)u = x(z(xu))$ holds.*

Proof. Let $g : A \times A \rightarrow F$ be a nondegenerate associative symmetric bilinear form, and let x, y, z, u be in A . Then we have

$$\begin{aligned} g\left(y((xz)x) - ((yx)z)x, u\right) &= g((xz)x, uy) - g(z, (xu)(yx)) \\ &= g\left(z, (x(uy))x - (xu)(yx)\right). \end{aligned} \tag{6}$$

On the other hand,

$$\begin{aligned} g\left(z, (x(uy))x - (xu)(yx)\right) &= g(xz, x(uy)) - g(z(xu), yx) \\ &= g\left(\left((xz)x\right)u - x(z(xu)), y\right). \end{aligned} \tag{7}$$

Since g is nondegenerate, (6) and (7) yield the equivalence of Assertions (i), (ii), and (iii). ■

THEOREM 1.2. *Let V be an H^* -algebra. Then the following assertions are equivalent:*

- (i) *V satisfies the right Moufang identity.*
- (ii) *V satisfies the middle Moufang identity.*
- (iii) *For all $x, z, u \in V$ the equality $((xz)x)u = x(z(xu))$ holds.*
- (iv) *V satisfies the left Moufang identity.*
- (v) *V is an alternative algebra.*

Proof. Let $g : V \times V \longrightarrow \mathbb{K}$ be the symmetric bilinear form defined by $g(x, y) = (x|y^*) + (y|x^*)$ for all $x, y \in V$. A routine calculation shows that g is associative and nondegenerate. Now Lemma 1.1 yields the equivalence of (i), (ii), and (iii).

Assume that the equivalent Conditions (i) and (iii) are fulfilled. By (i), for $x, y, z, t \in V$ we have

$$\begin{aligned} ((xy)x - x(yx) | zt) &= \left(z^*((xy)x) | t \right) - \left(z^*(x(yx)) | t \right) \\ &= \left(((z^*x)y)x | t \right) - \left(z^* | t(x(yx))^* \right) \\ &= \left(z^* | ((tx^*)y^*)x^* \right) - \left(z^* | ((tx^*)y^*)x^* \right) = 0. \end{aligned}$$

Without loss of generality we can assume that $\text{Ann } V = \{0\}$. Then, since the linear span of $\{zt : z, t \in V\}$ is dense in V (see (1)), and $(\cdot | \cdot)$ is nondegenerate, we obtain that $(xy)x = x(yx)$ for all $x, y \in V$. In this way we have proved that V is a flexible algebra. By (iii), V is left Moufang.

Now assume that Condition (iv) is fulfilled. Since the opposite algebras of left Moufang H^* -algebras are right Moufang H^* -algebras, and these last algebras are flexible, we obtain that V is also flexible. Then, applying again that V is left Moufang, we realize that Assertion (iii) holds for V .

Finally, we assume that V satisfies the equivalent Assertions (i)-(iv), and show that V is alternative. In view of (1), (2), and [9, Theorem 1], we can assume that V is topologically simple and complex. We already know that V is flexible, and that, consequently, for x in V the equality $x^2x = xx^2$ holds. On the other hand, by Assertion (ii), we have $x^2x^2 = (xx^2)x$ for every $x \in V$. Since there exists a nondegenerate associative symmetric bilinear form on V , it follows from [1] (see also [25, p. 39]) that V is a noncommutative Jordan H^* -algebra. Then, by [32, Theorems 2 and 4] and [2], either V is anticommutative or there exists an approximate unit for V . Assume that V is anticommutative. Then, by Assertion (i), we have $R_x^3 = 0$ for every x in V , where R_x stands for the operator of right multiplication by x on V . Therefore, if x is in V and satisfies $x^* = \varepsilon x$ for $\varepsilon \in \{+, -\}$, then for every y in V we have $0 = (R_x^4(y)|y) = \|R_x^2(y)\|^2$, and hence $0 = (R_x^2(y)|y) = \varepsilon \|R_x(y)\|^2$, so that $x \in \text{Ann } V = \{0\}$. Since $V = \text{Sym}(V, \star) \oplus \text{Sk}(V, \star)$, where $\text{Sym}(V, \star)$ (respectively, $\text{Sk}(V, \star)$) means the real vector subspace of all self-adjoint (respectively, skew-adjoint) elements of V , we deduce $V = 0$, which contradicts that V is topologically simple. Now assume that V has an approximate unit (say $\{e_\lambda\}_{\lambda \in \Lambda}$). Then, making $z = e_\lambda$ in (3) and $y = e_\lambda$ in (5), and taking limits in λ , the alternativeness of V follows. ■

As we said in the introduction, alternative H^* -algebras are well-understood. Indeed, *the topologically simple alternative nonassociative H^* -algebras are: the algebra of complex octonions, in the complex case; and the same algebra regarded as a real algebra, together with the two real octonions algebras, in the real case* (see [44, p. 148]). On the other hand every topologically simple associative H^* -algebra is isomorphic to the H^* -algebra $\mathcal{HS}(H)$ of all Hilbert-Schmidt operators on a suitable Hilbert space H , which is complex in the complex case, and either real, complex, or quaternionic in the real case (see [2] and [39]).

REFERENCES

- [1] ALBERT, A.A., Power associative rings, *Trans. Amer. Math. Soc.*, **64** (1948), 552–593.
- [2] AMBROSE, W., Structure theorems for a special class of Banach algebras, *Trans. Amer. Math. Soc.*, **57** (1945), 364–386.
- [3] BALACHANDRAN, V.K., Simple L^* -algebras of classical type, *Math. Ann.*, **180** (1969), 205–219.
- [4] BALACHANDRAN, V.K., Real L^* -algebras, *Indian J. Pure Appl. Math.*, **3** (1972), 1224–1246.
- [5] BALACHANDRAN, V.K., SWAMINATHAN, N., Real H^* -algebras, *J. Funct. Anal.*, **65** (1986), 64–75.
- [6] BENSEBAH, A., JV -algebras et JH^* -algèbres, *Canad. Math. Bull.*, **34** (1991), 447–455.
- [7] CABRERA, M., MARRACKCHI, A. EL, MARTÍNEZ, J., RODRÍGUEZ, A., An Allison-Kantor-Koecher-Tits construction for Lie H^* -algebras, *J. Algebra*, **164**(2) (1994), 361–408.
- [8] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., Malcev H^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **103** (1988), 463–471.
- [9] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., Nonassociative real H^* -algebras, *Publicacions Matemàtiques*, **32** (1988), 267–274.
- [10] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., A note on real H^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **105** (1989), 131–132.
- [11] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., Structurable H^* -algebras, *J. Algebra*, **147** (1992), 19–62.
- [12] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., Hilbert modules over H^* -algebras in relation with Hilbert ternary rings, in “Nonassociative Algebraic Models”, S. González and H. Ch. Myung (editors), Nova Science Publishers, New York, 1992, 33–44.
- [13] CABRERA, M., MARTÍNEZ, J., RODRÍGUEZ, A., Bounded differential operators on Hilbert modules and derivations of structurable H^* -algebras, *Comm. Algebra*, **21** (1993), 2905–2945.
- [14] CABRERA, M., RODRÍGUEZ, A., Extended centroid and central closure of semiprime normed algebras: a first approach, *Comm. Algebra*, **18** (1990), 2293–2326.

- [15] CABRERA, M., RODRÍGUEZ, A., Nonassociative ultraprime normed algebras, *Quart. J. Math. Oxford*, **43** (1992), 1–7.
- [16] CASTELLÓN, A., CUENCA, J.A., Compatibility in Jordan H^* -triple systems, *Bolletino U.M.I.*, **4**(B) (1990), 433–447.
- [17] CASTELLÓN, A., CUENCA, J.A., Alternative H^* -triple systems, *Comm. Algebra*, **11** (20) (1992), 3191–3260.
- [18] CASTELLÓN, A., CUENCA, J.A., Associative H^* -triple systems, in “Nonassociative Algebraic Models”, S. González and H. Ch. Myung (editors), Nova Science Publishers, New York, 1992, 45–67.
- [19] CASTELLÓN, A., CUENCA, J.A., Isomorphisms of H^* -triple systems, *Annali Della Scuola Normale Superiore di Pisa. Ser. IV*, **XIX** (4) (1992), 507–514.
- [20] CASTELLÓN, A., CUENCA, J.A., MARTÍN, C., Ternary H^* -algebras, *Bolletino U.M.I.*, **6-B** (7) (1992), 217–228.
- [21] CASTELLÓN, A., CUENCA, J.A., Centroid and metacentroid of an H^* -triple system, *Bulletin de la Societé Mathématique de Belgique*, **45** (fascicules 1-2 serie A) (1993), 85–91.
- [22] CASTELLÓN, A., CUENCA, J.A., MARTÍN, C., Applications of ternary H^* -algebras to associative H^* -superalgebras, *Algebras, Groups and Geometries*, **10** (1993), 181–190.
- [23] CASTELLÓN, A., CUENCA, J.A., MARTÍN, C., Jordan H^* -triple systems, in “Nonassociative Algebras and its Applications”, S. González (editor), Kluwer, Dordrecht-Boston-New York, 1994, 66–72.
- [24] CASTELLÓN, A., CUENCA, J.A., MARTÍN, C., Special Jordan H^* -triple systems, *Comm. Algebra*, **28** (10) (2000), 4699–4706.
- [25] CUENCA, J.A., Sobre H^* -álgebras no asociativas. Teoría de estructura de las H^* -álgebras de Jordan no conmutativas semisimples, Secretariado de Publicaciones de la Universidad de Málaga, Málaga, 1984.
- [26] CUENCA, J.A., Sur la theorie de structure des H^* -algèbres de Jordan non commutatives, *Ann. Sci. Univ. Clermont-Ferrand II Math.*, **27** (1991), 143–152.
- [27] CUENCA, J.A., On division algebras satisfying Moufang identities, preprint.
- [28] CUENCA, J.A., Composition algebras satisfying Moufang identities, preprint.
- [29] CUENCA, J.A., GARCÍA, A., MARTÍN, C., Structure theory for L^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **107** (1990), 361–365.
- [30] CUENCA, J.A., MARTÍN, C., Jordan two-graded H^* -álgebras, in “Nonassociative Algebraic Models”, S. González and H. Ch. Myung (editors), Nova Science Publishers, New York, 1992, 91–100.
- [31] CUENCA, J.A., RODRÍGUEZ, A., Isomorphisms of H^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **97** (1985), 93–99.
- [32] CUENCA, J.A., RODRÍGUEZ, A., Structure theory for noncommutative Jordan H^* -algebras, *J. Algebra*, **106** (1987), 1–14.
- [33] CUENCA, J.A., RODRÍGUEZ, A., Absolute values on H^* -algebras, *Comm. Algebra*, **23** (5) (1995), 2595–2610.
- [34] CUENCA, J.A., SÁNCHEZ, A., Structure theory for real noncommutative Jordan H^* -algebras, *J. Algebra*, **164** (1994), 481–499.

- [35] DE LA HARPE, P., Classification des L^* -algebres semi-simples réelles separables, *C.R. Acad. Sci, Ser A*, **272** (1971), 1559–1561.
- [36] DEVAPAKKIAM, C. VIOLA, Hilbert space methods in the theory of Jordan algebras I, *Math. Proc. Camb. Phil. Soc.*, **78** (1975), 293–300.
- [37] DEVAPAKKIAM, C. VIOLA, REMA, P.S., Hilbert space methods in the theory of Jordan algebras II, *Math. Proc. Camb. Phil. Soc.*, **79** (1976), 307–319.
- [38] FERNÁNDEZ, A., RODRÍGUEZ, A., A Wedderburn theorem for nonassociative complete normed algebras, *J. London Math. Soc.*, **33** (1986), 328–338.
- [39] KAPLANSKY, I., Dual rings, *Ann. of Math.*, **49** (3) (1948), 689–701.
- [40] KAUP, W., Über die klassifikation der symmetrischen hermiteschen mannigfaltigkeiten unendlicher dimension II, *Math. Ann.*, **262** (1983), 57–75.
- [41] NEHER, E., “Jordan Triple Systems by the Grid Approach”, Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [42] NEHER, E., Generators and relations for 3-graded Lie algebras, *J. Algebra*, **155** (1) (1993), 1–35.
- [43] PÉREZ DE GUZMÁN, I., Structure theorems for alternative H^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **94** (1983), 437–446.
- [44] RODRÍGUEZ, A., Jordan structures in Analysis, in “Jordan Algebras”, K. McCrimmon, W. Kaup, and H. P. Petersson (editors), Walter de Gruyter, Berlin, 1994, 97–186.
- [45] RODRÍGUEZ, A., Continuity of densely valued homomorphisms into H^* -algebras, *Quart. J. Math. Oxford*, **46** (1995), 107–118.
- [46] SCHAFER, R.D., “An Introduction to Nonassociative Algebras”, Academic Press, New York-San Francisco-London, 1966.
- [47] SCHUE, J.R., Hilbert space methods in the theory of Lie algebras, *Trans. Amer. Math. Soc.*, **95** (1960), 68–80.
- [48] SCHUE, J.R., Cartan decompositions for L^* -algebras, *Trans. Amer. Math. Soc.*, **98** (1961), 334–349.
- [49] UNSAIN, I., Classification of the simple separable real L^* -algebras, *J. Differential Geom.*, **7** (106) (1972), 423–451.
- [50] VILLENA, A., Continuity of derivations on H^* -algebras, *Proc. Amer. Math. Soc.*, **122** (1994), 821–826.
- [51] VILLENA, A.R., ZALAR, B., ZOHRY, M., Continuity of derivations pairs on Hilbert triple systems, *Bollettino U.M.I.*, **7** (9-B) (1995), 459–477.
- [52] VILLENA, A., ZHORY, M., Continuity of homomorphism pairs into H^* -triple systems, *Proc. (Math. Sci.) I. A. Sc.*, **106** (1996), 79–90.
- [53] VILLENA, A., ZHORY, M., Rings isomorphisms of H^* -algebras, *J. Algebra*, **181** (1996), 329-346.
- [54] VILLENA, A., ZHORY, M., Rings derivations on H^* -algebras, *Bollettino U.M.I.*, **11-A** (1997), 129–138.
- [55] ZHEVLAKOV, K.A., SLIN’KO, A.M., SHESTAKOV, I.P., SHIRSHOV, A.I., “Rings that are Nearly Associative”, Academic Press, New York, 1982.
- [56] ZALAR, B., Continuity of derivations on Malcev H^* -algebras, *Math. Proc. Camb. Phil. Soc.*, **110** (1991), 455–459.

- [57] ZALAR, B., Continuity of derivations pairs on alternative and Jordan H^* -triples, *Boll. Un. Mat. Ital.*, **7** (B) (1993), 215–230.
- [58] ZALAR, B., Theory of Hilbert triple systems, *Yokohama Math. J.*, **41** (1994), 95–126.